

A Note on the Radiatively Induced Lorentz and CPT Violated Chern-Simons-like Term in the Extended Quantum Electrodynamics

W.F. Chen*

*Department of Mathematics,
Nipissing University, North Bay,
Ontario, Canada P1B 8L7*

and

*Department of Physics,
University of Winnipeg, Winnipeg,
Manitoba, Canada R3B 2E9*

We show that the ambiguity for the Chern-Simons-like term induced from quantum correction in the extended QED should have nothing to do with the approximation on the exact fermionic propagator, contradictory to the claim in Ref.[19]. Further, we investigate the induced Chern-Simons-like term using the original 't Hooft-Veltman dimensional regularization and reproduce the result obtained by gauge symmetry analysis. This fact demonstrates that the origin of the ambiguity should lie in different choices on regularization schemes.

PACS numbers: 11.15.Bt, 11.30.Cp, 12.20.Ds, 11.40.Ha

It is more than tens years that the radiatively induced Chern-Simons-like term in an extended QED has been keeping as a popular topic. The most remarkable feature of this topic is the ambiguity for the induced Chern-Simons-like term: a number of distinct results have been obtained [1]–[19]. As pointed out by Jackiw [20], this ambiguity is a typical feature of quantum field theory when some fundamental symmetry is violated that “radiative corrections are finite but undetermined”. A convincing viewpoint is that the ambiguity originates from the different regularization schemes of tackling the linear UV divergence [5]. However, recently it has been proposed that the ambiguity may originate from different approximations on the exact fermionic propagator and should be independent of the utilization on the regularization methods [19]. In this letter we show explicitly that the claimed difference in the approximation on the exact fermionic propagator actu-

*Electronic address: wengfenc@nipissingu.ca

ally does not exist. Further, we show that the origin for the discrepancy in the radiatively induced Chern-Simons like term should attribute to the choices on regularization schemes.

The fermionic sector of the extended quantum electrodynamics containing a Lorentz- and CPT-violating axial vector interaction with a constant four-vector is described by the following Lagrangian density:

$$\mathcal{L}_F = \bar{\psi} (i\partial - e\mathcal{A} - m - \gamma_5 \not{b}) \psi. \quad (1)$$

The exact fermionic propagators with the constant vector field b_μ reads

$$S(p, b) = \frac{i}{\not{p} - m - \not{b}\gamma_5} \quad (2)$$

In the following we show that the two different approximations on the propagator (2) claimed in Ref. [19] are actually identical. First, the approximation expansion of (2) to the linear order of b_μ given in Ref. [14] is

$$\begin{aligned} S_I(p, b) &= \frac{i(\not{p} + m + \not{b}\gamma_5)}{p^2 - m^2 + b^2 - 2m\not{b}\gamma_5 + p \cdot b\gamma_5} = \frac{i(\not{p} + m + \not{b}\gamma_5)}{p^2 - m^2} \left[1 + \frac{2m\not{b} - 2p \cdot b\gamma_5}{p^2 - m^2} \right] + \mathcal{O}(b^2) \\ &= \frac{i(\not{p} + m + \not{b}\gamma_5)}{p^2 - m^2} - \frac{2i\gamma_5(m\not{b} - p \cdot b)(\not{p} + m^2)}{(p^2 - m^2)^2} + \mathcal{O}(b^2) \\ &= \frac{i(\not{p} + m + \not{b}\gamma_5)}{p^2 - m^2} - \frac{2i\gamma_5(mb \cdot p - \not{p}b \cdot p - m\not{p}\not{b} + m^2\not{b})}{(p^2 - m^2)^2} + \mathcal{O}(b^2) \end{aligned} \quad (3)$$

On the other hand, with the identity

$$\frac{1}{A+B} = \frac{1}{A} - \frac{1}{A}B\frac{1}{A+B} = \frac{1}{A} - \frac{1}{A}B\left(\frac{1}{A} - \frac{1}{A}B\frac{1}{A+B}\right), \quad (4)$$

there arises another approximation expansion of Eq.(2), which has been used in most of the literatures,

$$\begin{aligned} S_{II}(p, b) &= \frac{i}{\not{p} - m} + i\frac{1}{\not{p} - m}\not{b}\gamma_5\frac{1}{\not{p} - m} + \mathcal{O}(b^2) \\ &= \frac{i(\not{p} + m)}{p^2 - m^2} + \frac{i(\not{p} + m)\not{b}\gamma_5(\not{p} + m)}{(p^2 - m^2)^2} \\ &= \frac{i(\not{p} + m + \not{b}\gamma_5)}{p^2 - m^2} + \frac{i\gamma_5(2p \cdot b\not{p} - 2m^2\not{b} + m\not{p}\not{b} - m\not{b}\not{p})}{(p^2 - m^2)^2} + \mathcal{O}(b^2) \\ &= \frac{i(\not{p} + m + \not{b}\gamma_5)}{p^2 - m^2} - \frac{2i\gamma_5(mb \cdot p - \not{p}b \cdot p - m\not{p}\not{b} + m^2\not{b})}{(p^2 - m^2)^2} + \mathcal{O}(b^2) \end{aligned} \quad (5)$$

One can see from Eqs. (3) and (5) that two approximation expansions are identical: $S_I(p, b) = S_{II}(p, b)$.

The above expansions are performed with no account taken on the regularization. To be more confirmative, we consider the 't Hooft-Veltman dimensional regularization [21, 22] and observe whether there exists difference between two approximation approaches at the regularization level.

The prescription for the 't Hooft-Veltman is defined as follows [21, 22]: the regularized n -dimensional space is divided into a direct sum of the original four-dimensional and $n-4$ -dimensional spaces. When one calculates the amplitude represented by an Feynman diagram, the external momentum p_μ lives only in the four-dimensional space, and the loop momentum k_μ in the whole n -dimensional space,

$$\begin{aligned} g_{\mu\nu} &= \tilde{g}_{\mu\nu} \oplus \hat{g}_{\mu\nu}, & p_\mu &= \tilde{p}_\mu \oplus 0 & k_\mu &= \tilde{k}_\mu \oplus \hat{k}_\mu, \\ \tilde{g}_{\mu\nu} \hat{k}^\nu &= \hat{g}_{\mu\nu} \tilde{k}^\nu = 0, & \tilde{g}_{\mu\nu} &= \eta_{\mu\nu}, & \hat{g}_{\mu\nu} &= -\delta_{\mu\nu}. \end{aligned} \quad (6)$$

The $2^{[n/2]} \times 2^{[n/2]}$ γ -matrices are defined as

$$\begin{aligned} \gamma_\mu &= (\tilde{\gamma}_\mu, \hat{\gamma}_\mu), & \gamma_5 &= \gamma^5 \equiv i\tilde{\gamma}^0\tilde{\gamma}^1\tilde{\gamma}^2\tilde{\gamma}^3 = \frac{i}{4!}\epsilon_{\mu\nu\lambda\rho}\tilde{\gamma}^\mu\tilde{\gamma}^\nu\tilde{\gamma}^\lambda\tilde{\gamma}^\rho, \\ \{\gamma_\mu, \gamma_\nu\} &= 2g_{\mu\nu}, & \{\tilde{\gamma}_\mu, \tilde{\gamma}_\nu\} &= 2\tilde{g}_{\mu\nu}, & \{\hat{\gamma}_\mu, \hat{\gamma}_\nu\} &= 2\hat{g}_{\mu\nu}, \\ \{\tilde{\gamma}_\mu, \hat{\gamma}_\nu\} &= 0, & \{\gamma_5, \tilde{\gamma}_\mu\} &= 0, & [\gamma_5, \hat{\gamma}_\mu] &= 0, \end{aligned} \quad (7)$$

where $\tilde{\gamma}_\mu$ has only non-vanishing 4×4 components in the upper-left corner and $\hat{\gamma}_\mu$ has only the non-vanishing $2^{-\epsilon} \times 2^{-\epsilon}$ elements in the lower-right corner, $\epsilon \equiv 2 - n/2$. In addition, like the external momentum, the γ -matrix appearing in the external vertex lives in four dimensions. One can easily see that the above regularization prescription presents the $SO(1,3) \times SO(n-3)$ covariance rather than the whole $SO(n)$ covariance.

According to the above prescription, the first approximation expansion on the exact fermionic propagator appearing in a loop of a certain Feynman diagram with a loop momentum k is

$$\begin{aligned} S_I^R(k, b) &= \frac{i}{\tilde{k}^2 + \hat{k}^2 - m - \not{b}\gamma_5} = \frac{i(\tilde{k} + \hat{k} + m + \not{b}\gamma_5)}{\tilde{k}^2 - \hat{k}^2 - m^2 + b^2 - 2m\not{b}\gamma_5 + 2\tilde{k} \cdot b\gamma_5 - 2\not{b}\hat{k}\gamma_5} \\ &= \frac{i(\tilde{k} + \hat{k} + m + \not{b}\gamma_5)}{\tilde{k}^2 - \hat{k}^2 - m^2} - \frac{2i\gamma_5(m\not{b} - \tilde{k} \cdot b)(\tilde{k} + \hat{k} + m)}{(\tilde{k}^2 - \hat{k}^2 - m^2)^2} \\ &\quad + \frac{2i\gamma_5\not{b}(m\hat{k} - \hat{k}^2 - \tilde{k}\hat{k})}{(\tilde{k}^2 - \hat{k}^2 - m^2)^2} + \mathcal{O}(b^2) \end{aligned} \quad (8)$$

While the second approximation leads to

$$\begin{aligned} S_{II}^R(k, b) &= \frac{i(\tilde{k} + \hat{k} + m)}{\tilde{k}^2 - \hat{k}^2 - m^2} + \frac{i(\tilde{k} + \hat{k} + m)\not{b}\gamma_5(\tilde{k} + \hat{k} + m)}{(\tilde{k}^2 - \hat{k}^2 - m^2)^2} + \mathcal{O}(b^2) \\ &= \frac{i(\tilde{k} + \hat{k} + m + \not{b}\gamma_5)}{\tilde{k}^2 - \hat{k}^2 - m^2} - \frac{2i\gamma_5(m\not{b} - \tilde{k} \cdot b)(\tilde{k} + \hat{k} + m)}{(\tilde{k}^2 - \hat{k}^2 - m^2)^2} \end{aligned}$$

$$+ \frac{2i\gamma_5 \not{b} (m\hat{k} - \hat{k}^2 - \tilde{k}\tilde{k})}{(\tilde{k}^2 - \hat{k}^2 - m^2)^2} + \mathcal{O}(b^2) \quad (9)$$

Eqs. (8) and (9) show that even at the 't Hooft-Veltman dimensional regularization level, the two b -linear order approximations for the exact fermionic propagator appearing in the fermionic loop are still identical:

$$S_I^R(k, b) = S_{II}^R(k, b) \equiv S^R(k, b) \quad (10)$$

In the following we observe the induced Chern-Simons-like terms using the prescription defined in (6) and (7). The one-loop vacuum polarization tensor in the 't Hooft-Veltman dimensional regularization is

$$\begin{aligned} i\Pi_{\mu\nu}^{(1)R}(p, b) &= - \int \frac{d^n k}{(2\pi)^n} \text{Tr} \left[(-ie\mu^\epsilon \gamma_\mu) S^R(k) (-ie\mu^\epsilon \gamma_\nu) S^R(k+p) \right] \\ &= -e^2 \mu^{2\epsilon} \int \frac{d^n k}{(2\pi)^n} \text{Tr} \left(\gamma_\mu \frac{1}{\not{k} - m - \not{b}\gamma_5} \gamma_\nu \frac{1}{\not{k} + \not{p} - \not{b}\gamma_5} \right) \end{aligned} \quad (11)$$

Using the expansion (4), we focus on the sector of the vacuum polarization tensor relevant to the Chern-Simons-like term, which consists of the b -linear terms in $\Pi_{\mu\nu}^{(1)}(p, b)$,

$$\begin{aligned} i\Pi_{\mu\nu}^{\text{CS}}(p, b) &= i\Pi_{\mu\nu}^{(1)\text{CS}}(p, b) + i\Pi_{\mu\nu}^{(2)\text{CS}}(p, b), \\ i\Pi_{\mu\nu}^{(1)\text{CS}}(p, b) &= -e^2 \mu^{2\epsilon} \int \frac{d^n k}{(2\pi)^n} \text{Tr} \left(\gamma_\mu \frac{1}{\not{k} - m} \not{b}\gamma_5 \frac{1}{\not{k} - m} \gamma_\nu \frac{1}{\not{k} + \not{p} - m} \right), \\ i\Pi_{\mu\nu}^{(2)\text{CS}}(p, b) &= -e^2 \mu^{2\epsilon} \int \frac{d^n k}{(2\pi)^n} \text{Tr} \left(\gamma_\mu \frac{1}{\not{k} - m} \gamma_\nu \frac{1}{\not{k} + \not{p} - m} \not{b}\gamma_5 \frac{1}{\not{k} + \not{p} - m} \right). \end{aligned} \quad (12)$$

Further, one can (naively) show that $\Pi_{\mu\nu}^{(1)\text{CS}}(p, b)$ and $\Pi_{\mu\nu}^{(2)\text{CS}}(p, b)$ are identical. Making the shift $k \rightarrow k - p$ in $\Pi_{\mu\nu}^{(2)\text{CS}}(p, b)$, we have

$$\begin{aligned} i\Pi_{\mu\nu}^{(2)\text{CS}}(p, b) &= -e^2 \mu^{2\epsilon} \int \frac{d^n k}{(2\pi)^n} \text{Tr} \left(\gamma_\mu \frac{1}{\not{k} - \not{p} - m} \gamma_\nu \frac{1}{\not{k} - m} \not{b}\gamma_5 \frac{1}{\not{k} - m} \right) \\ &= -e^2 \mu^{2\epsilon} \int \frac{d^n k}{(2\pi)^n} \text{Tr} \left(\gamma_\nu \frac{1}{\not{k} - m} \not{b}\gamma_5 \frac{1}{\not{k} - m} \gamma_\mu \frac{1}{\not{k} - \not{p} - m} \right) \\ &= i\Pi_{\nu\mu}^{(1)\text{CS}}(-p, b) = \Pi_{\mu\nu}^{(1)\text{CS}}(p, b) \end{aligned} \quad (13)$$

where in the last line we have used the tensor structure of $\Pi_{\mu\nu}^{\text{CS}}(p, b) = \epsilon_{\mu\nu\lambda\rho} p^\lambda b^\rho f(p^2)$. Therefore, we obtain

$$\begin{aligned} i\Pi_{\mu\nu}^{\text{CS}}(p, b) &= 2i\Pi_{\mu\nu}^{(1)\text{CS}}(p, b) \\ &= -2e^2 \mu^{2\epsilon} \int \frac{d^n k}{(2\pi)^n} \frac{\text{Tr} \left[\gamma_\mu (\not{k} + \not{k} + m) \not{b}\gamma_5 (\not{k} + \not{k} + m) \gamma_\nu (\not{k} + \not{p} + \not{k} + m) \right]}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} \\ &= 8ie^2 \mu^{2\epsilon} \epsilon_{\mu\nu\lambda\rho} \int \frac{d^4 \tilde{k}}{(2\pi)^4} \int \frac{d^{n-4} \hat{k}}{(2\pi)^{n-4}} \frac{2\tilde{k} \cdot b \tilde{k}^\lambda p^\rho + (\tilde{k}^2 - \hat{k}^2 - m^2) \tilde{k}^\lambda b^\rho + (\tilde{k}^2 + \hat{k}^2 + m^2) p^\lambda b^\rho}{(\tilde{k}^2 - \hat{k}^2 - m^2)^2 [(\tilde{k} + p)^2 - \hat{k}^2 - m^2]} \end{aligned} \quad (14)$$

In above we have used the prescriptions listed in (6) and (7) and the following formula,

$$\text{Tr}(\gamma_5 \tilde{\gamma}_\mu \tilde{\gamma}_\nu \tilde{\gamma}_\lambda \tilde{\gamma}_\rho) = -4i\epsilon_{\mu\nu\lambda\rho}, \quad \widehat{k}\widehat{k} = -\widehat{k}^2 \quad (15)$$

To evaluate the non-covariant integrals in Eq.(14), it is very convenient to repeatedly use the following decomposition to improve the UV behavior of the integrand,

$$\frac{1}{(k+p)^2 - m^2} = \frac{1}{k^2 - m^2} - \frac{2k \cdot p + p^2}{(k^2 - m^2)[(k+p)^2 - m^2]}. \quad (16)$$

Once the integration becomes UV finite, one can directly take the $n \rightarrow 4$ limit before performing the integration, and the integrand proportional to the evanescent quantity $\widehat{k}_\mu$ will vanish automatically [23]. For examples, we have

$$\begin{aligned} & \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\widehat{k}^2}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} \\ &= \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\widehat{k}^2}{(k^2 - m^2)^2} \left(\frac{1}{k^2 - m^2} - \frac{2k \cdot p + p^2}{(k^2 - m^2)[(k+p)^2 - m^2]} \right) \\ &= \lim_{n \rightarrow 4} \frac{1}{(2\pi)^n} \int d^4 \tilde{k} \int d^{n-4} \widehat{k} \frac{\widehat{k}^2}{(\tilde{k}^2 - \widehat{k}^2 - m^2)^3} = \frac{i}{32\pi^2}; \end{aligned} \quad (17)$$

$$\begin{aligned} & \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\tilde{k}_\mu \tilde{k}_\nu}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} \\ &= \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\tilde{k}_\mu \tilde{k}_\nu}{(k^2 - m^2)^2} \left(\frac{1}{k^2 - m^2} - \frac{2\tilde{k} \cdot p + p^2}{(k^2 - m^2)[(k+p)^2 - m^2]} \right) \\ &= \frac{1}{4} \tilde{g}_{\mu\nu} \lim_{n \rightarrow 4} \int \frac{d^4 \tilde{k}}{(2\pi)^4} \tilde{k}^2 \int \frac{d^{n-4} \widehat{k}}{(2\pi)^{n-4}} \frac{1}{(\tilde{k}^2 - \widehat{k}^2 - m^2)^3} - \int \frac{d^4 k}{(2\pi)^4} \frac{(2k \cdot p + p^2) k_\mu k_\nu}{(k^2 - m^2)^3 [(k+p)^2 - m^2]} \\ &= \frac{i}{64\pi^2} g_{\mu\nu} \left[\frac{1}{\epsilon} - \gamma + \ln \frac{4\pi\mu^2}{m^2} + 2 - \frac{2m}{p} \left(4 - \frac{p^2}{m^2} \right)^{1/2} \arctan \frac{p/m}{(4 - p^2/m^2)^{1/2}} \right] \\ &+ \frac{i}{32\pi^2} \frac{p_\mu p_\nu}{p^2} \left[1 - 4 \frac{m}{p} \frac{1}{(4 - p^2/m^2)^{1/2}} \arctan \frac{p/m}{(4 - p^2/m^2)^{1/2}} \right] \end{aligned} \quad (18)$$

Note that in above integrations one should integrate over $\widehat{k}$ first and then over $\tilde{k}$.

Using the integration formula listed in Appendix, we can obtain the part of the vacuum polarization tensor relevant to the Chern-Simon-like term,

$$\begin{aligned} i\Pi_{\mu\nu}^{\text{CS}}(p, b) &= \frac{ie^2}{2\pi^2} \epsilon_{\mu\nu\lambda\rho} p^\lambda b^\rho \left[\frac{1}{2} - \int_0^1 dx \frac{1-x}{1-x(1-x)p^2/m^2} \right. \\ &\quad \left. + \frac{p^2}{m^2} \int_0^1 dx \frac{5x(1-x)^2 - 2(1-x)^3 - x(1-x)}{1-x(1-x)p^2/m^2} \right] \\ &= \frac{ie^2}{2\pi^2} \epsilon_{\mu\nu\lambda\rho} p^\lambda b^\rho \left[-4 + \frac{4m}{p} \left(4 - \frac{p^2}{m^2} \right)^{1/2} \arctan \frac{p/m}{(4 - p^2/m^2)^{1/2}} \right] \end{aligned} \quad (19)$$

The induced Chern-Simons coefficient k_μ is defined from $\Pi_{\mu\nu}^{\text{CS}}(p, b)$ at $p^2 = 0$. Eq. (19) gives

$$k_\mu = 0 \quad (20)$$

However, this is not the end of story, one should evaluate the surface term produced by the loop momentum shift $k \rightarrow k - p$, since the loop integral is linearly divergent. As shown in Ref. [2], the shift can somehow lead to a non-vanishing surface term. In spite of the overwhelming viewpoint that the shift can be safely taken after having implemented a regularization, but due to the particularity of the t'Hooft-Veltman regularization scheme, an explicit calculation should be performed to observe whether a finite result can arise from the surface term. The calculation on the chiral anomaly from the triangle diagram is a typical example [24].

The surface term induced by the variable shift in Eq. (13) is

$$\begin{aligned} \Delta\Pi_{\mu\nu}^{(2)CS}(p, b) = & ie^2\mu^{2\epsilon} \int \frac{d^n k}{(2\pi)^n} \left[\text{Tr} \left(\gamma_\mu \frac{1}{\not{k} - m} \gamma_\nu \frac{1}{\not{k} + \not{p} - m} \not{p} \gamma_5 \frac{1}{\not{k} + \not{p} - m} \right) \right. \\ & \left. - \text{Tr} \left(\gamma_\mu \frac{1}{\not{k} - \not{p} - m} \gamma_\nu \frac{1}{\not{k} - m} \not{p} \gamma_5 \frac{1}{\not{k} - m} \right) \right] \end{aligned} \quad (21)$$

The calculation on the surface term is the same as that evaluating the chiral anomaly from the triangle diagram $\langle j_\mu(p) j_\nu(p - q) j_\rho^5(q) \rangle$ with the momentum of the axial vector current $q = 0$. In the 't Hooft-Veltman dimensional regularization, the surface term can only get contribution from the integrands containing the evanescent momentum $\hat{k}^2$ [24]. Therefore, evaluating the trace in (21) and taking into account the terms containing $\hat{k}^2$, we have

$$\begin{aligned} \Delta\Pi_{\mu\nu}^{(2)CS}(p, b) = & 4e^2 \epsilon_{\mu\nu\lambda\rho} \int \frac{d^n k}{(2\pi)^n} \left[\frac{\hat{k}^2 (\tilde{k}^\lambda + 2p^\lambda) b^\rho}{(k^2 - m^2)[(k + p)^2 - m^2]^2} - \frac{\hat{k}^2 (\tilde{k}^\lambda + p^\lambda) b^\rho}{(k^2 - m^2)^2[(k - p)^2 - m^2]} \right] \\ = & \frac{ie^2}{24\pi^2} \epsilon_{\mu\nu\lambda\rho} p^\lambda b^\rho (-2 + 6 - 1 - 3) = 0. \end{aligned} \quad (22)$$

This coincides with the result obtained in Ref. [9].

In summary, we have shown the two approximations on the exact fermionic propagator used in Ref. [19] are actually identical at both unregularized and regularization level. The discrepancy shown by the author should come from the different algebraic operations the author made on the integrand rather than taking approximation on the fermionic propagator. Further, using the 't Hooft-Veltman dimensional regularization, we have calculated the induced Chern-Simons-like term and obtain a vanishing result. The reason for this result is probably because the dimensional regulation preserves the Ward identities and make the theory well defined at regularization level, as analyzed in Ref. [9]. In comparison with the various results obtained in the literature [1]– [19],

the origin for the ambiguity in the induced Chern-Simons terms should lie in different choices on regularization schemes.

Acknowledgement: This work is supported by the Natural Sciences and Engineering Research Council of Canada. I would like to thank Professor R. Jackiw for encouragement. I also would like to thank Professor G. Bonneau for continuous comments.

APPENDIX A: INTEGRATION FORMULA

Some integration formula used to evaluate (14) and (21) are listed in this appendix:

$$\begin{aligned} \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\hat{k}^2}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} &= \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\hat{k}^2}{(k^2 - m^2) [(k+p)^2 - m^2]^2} \\ &= \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\hat{k}^2}{(k^2 - m^2)^3} = \lim_{n \rightarrow 4} \frac{1}{(2\pi)^n} \int d^4 \tilde{k} \int d^{n-4} \hat{k} \frac{\hat{k}^2}{(\tilde{k}^2 - \hat{k}^2 - m^2)^3} = \frac{i}{32\pi^2}, \end{aligned} \quad (A1)$$

$$\begin{aligned} \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\hat{k}^2 \tilde{k}_\mu}{(k^2 - m^2)^2 [(k-p)^2 - m^2]} &= \int \frac{d^n k}{(2\pi)^n} \frac{2\hat{k}^2 \tilde{k} \cdot p \tilde{k}_\mu}{(k^2 - m^2)^4} \\ &= \lim_{n \rightarrow 4} \frac{1}{(2\pi)^n} \int d^4 \tilde{k} \int d^{n-4} \hat{k} \frac{(-2)\hat{k}^2 \tilde{k}_\mu}{(\tilde{k}^2 - \hat{k}^2 - m^2)^4} = \frac{i}{96\pi^2} p_\mu, \end{aligned} \quad (A2)$$

$$\lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\hat{k}^2 \tilde{k}_\mu}{(k^2 - m^2) [(k+p)^2 - m^2]^2} = \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{(-4)\tilde{k}_\mu \hat{k}^2 \tilde{k} \cdot p}{(k^2 - m^2)^3} = -\frac{i}{48\pi^2} p_\mu \quad (A3)$$

$$\int \frac{d^4 k}{(2\pi)^4} \frac{1}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} = -\frac{i}{16\pi^2} \int_0^1 dx \frac{1-x}{1-x(1-x)p^2/m^2}, \quad (A4)$$

$$\int \frac{d^4 k}{(2\pi)^4} \frac{\tilde{k}_\mu}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} = \frac{i}{16\pi^2} p_\mu \frac{1}{m^2} \int_0^1 dx \frac{x(1-x)}{1-x(1-x)p^2/m^2}, \quad (A5)$$

$$\begin{aligned} \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\tilde{k}_\mu \tilde{k}_\nu}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} &= \frac{i}{64\pi^2} g_{\mu\nu} \left(\frac{1}{\epsilon} - \gamma + \ln \frac{4\pi\mu^2}{m^2} \right) \\ &+ \frac{i}{64\pi^2} g_{\mu\nu} \frac{p^2}{m^2} \int_0^1 dx \frac{(1-x)^3 - x(1-x)^2}{1-x(1-x)p^2/m^2} - \frac{i}{16\pi^2} \frac{p_\mu p_\nu}{p^2} \frac{p^2}{m^2} \int_0^1 dx \frac{x(1-x)^2}{1-x(1-x)p^2/m^2}, \end{aligned} \quad (A6)$$

$$\begin{aligned} \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\tilde{k}^2 \tilde{k}_\mu}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} &= \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\tilde{k}_\mu}{(k^2 - m^2) [(k+p)^2 - m^2]} \\ &+ \lim_{n \rightarrow 4} \int \frac{d^n k}{(2\pi)^n} \frac{\tilde{k}_\mu \hat{k}^2}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} + m^2 \int \frac{d^4 k}{(2\pi)^4} \frac{\tilde{k}_\mu}{(k^2 - m^2)^2 [(k+p)^2 - m^2]} \\ &= -\frac{i}{32\pi^2} p_\mu \left(\frac{1}{\epsilon} - \gamma + \ln \frac{4\pi\mu^2}{m^2} \right) - \frac{i}{32\pi^2} p_\mu \frac{p^2}{m^2} \int_0^1 dx \frac{(1-x)^3 - 5x(1-x)^2 + 2x(1-x)}{1-x(1-x)p^2/m^2} \\ &- \frac{i}{96\pi^2} p_\mu + \frac{i}{16\pi^2} p_\mu \int_0^1 dx \frac{x(1-x)}{1-x(1-x)p^2/m^2}, \end{aligned} \quad (A7)$$

$$(A8)$$

-
- [1] S. Coleman and S.L. Glashow, Phys. Rev. **D59** (1999) 116008.
 - [2] R. Jackiw and V.A. Kostelecký, Phys. Rev. Lett. **82** (1999) 3572.
 - [3] J.M. Chung and P. Oh, Phys. Rev. **D60** (1999) 067702; J.M. Chung, Phys. Rev. **D60** (1999) 127901; J.M. Chung, Phys. Lett. **B461** (1999) 138; J.M. Chung and B.K. Chung, Phys. Rev. **D63** (2001) 105015.
 - [4] W.F. Chen, Phys. Rev. **D60** (1999) 085007; M. Chaichian, W.F. Chen and R. Gonzalez Felipe, Phys. Lett. **B503** (2001) 215; W.F. Chen and G. Kunstatter, Phys. Rev. **D62** (2000) 105029.
 - [5] W.F. Chen, *Issues on radiatively induced Lorentz and CPT violation in quantum electrodynamics*, hep-th/0106035.
 - [6] M. Pérez-Victoria, Phys. Rev. Lett. **83** (1999) 2518; JHEP **0104** (2001) 032.
 - [7] L.-H. Chan, *Induced Lorentz-violating Chern-Simons term in QED and anomalous contributions to effective action expansions*, hep-ph/9907349.
 - [8] C. Adam and F.R. Klinkhamer, Nucl. Phys. **B607** (2001) 247; Phys. Lett. **B513** (2001) 245.
 - [9] G. Bonneau, Nucl. Phys. **B593** (2001) 398; Nucl. Phys. **B764** (2007) 83.
 - [10] A.P. Baeta Scarpelli, M. Sampaio, B. Hiller and M.C. Nemes, Phys. Rev. **D64** (2001) 046013; A. P. Baeta Scarpelli, H. Belich, J. L. Boldo and J. A. Helayel-Neto, Phys. Rev. **D67** (2003) 085021.
 - [11] Yu.A. Sitenko, Phys. Lett. **B515** (2001) 414; Yu.A. Sitenko and K.Yu. Rulik, Eur. Phys. J. **C28** (2005) 405.
 - [12] O.A. Battistel and G. Dallabona, Nucl. Phys. **B610**(2001) 316; Phys. Rev. **D72**(2005) 045009.
 - [13] A.A. Andrianov, P. Giacconi and R. Soldati, JHEP **0202** (2002) 030. J. Alfaro, A.A. Andrianov, M. Cambiaso, P. Giacconi and R. Soldati, Phys. Lett. **B639** (2006) 586.
 - [14] D. Ebert, V.Ch. Zhukovsky and A.S. Razumovsky, Phys. Rev. **D70** (2004) 025003.
 - [15] T. Mariz, J.R. Nascimento, E. Passos and R.F. Ribeiro, Phys. Rev. **D70** (2004) 024014; M. Gomes, J.R. Nascimento, E. Passos, A.Yu. Petrov and A.J. da Silva, Phys. Rev. **D76** (2007) 047701; T. Mariz, J.R. Nascimento, A.Yu. Petrov, L.Y. Santos and A.J. da Silva, arXiv: 0708.3348; M. Gomes, T. Mariz, J.R. Nascimento and A.J. da Silva, arXiv:0709.2904; F.A. Brito, J.R. Nascimento, E. Passos and A.Yu. Petrov, arXiv: 0709.3090.
 - [16] B. Altschul, Phys. Rev. **D70** (2004) 101701; *ibid* (2006) 045004; *ibid* (2006) 036005.
 - [17] Y.L. Ma and Y.L. Wu, Phys. Lett. **B647** (2007) 427.
 - [18] D. Colladay and P. McDonald, Phys. Rev. **D75** (2007) 105002.
 - [19] E. Passos, *Radiatively induced finite and (un)determined Chern-Simons-like terms*, arXiv: 0711.3992.
 - [20] R. Jackiw, Rajaramanfest lecture, New Delhi, March 1999, hep-th/9903044.
 - [21] G. 't Hooft and M. Veltman, Nucl. Phys. **B44**(1972) 189.
 - [22] P. Breitenlohner and D. Maison, Commu. Math. Phys. **52** (1977) 11.
 - [23] G. Giavarini, C.P. Martin and F. Ruiz Ruiz, Nucl. Phys. **B381** (1992) 222.

- [24] R.A. Bertlmann, *Anomalies in Quantum Field Theory*, Chapter 4, Oxford, 1996.